

Analysis of Reactive Processes in a CO₂ Pilot Injection Test


Aaron Abel, Brian McPherson,

Peter Lichtner, Rajesh Pawar, Norman Warpinski,
Gillian Bond, Ning Liu, John Stringer, and Reid Grigg

Outline


- Background
- Self-Trapping Concept
- Objectives
- Dawsonite Precipitation Timing study
- Conclusions
- Acknowledgements

West Pearl Queen Reservoir



New Mexico

Texas



Bulk Mineralogy

67% Quartz

23% Albite

5% K-Feldspar

2% Dolomite

2% Anhydrite

Strata Production Co. Wells

- #4 CO₂ Injection & monitoring well
- #5 Monitoring & producing well
- #1&3 Waste water injector well
- #2 Plugged well

Brine composition:

180,000 mg/L (NaCl-dominated)





Figure compliments of R. Pawar, LANL

Self-Trapping Concept

West Pearl Queen Reservoir: Shattuck Sandstone



Dawsonite

- Naturally occurring throughout regions of the world with CO₂-rich waters, but has not been reproduced within laboratory experiments

Specifically.....

- Springerville Natural CO₂ Field (eastern AZ)
(Moore et al., 2003)
- Bowen-Gunnedah-Sydney Basin (Australia)
occurs as a widespread cement within sedimentary rocks (Baker et al., 1995)
- Ordovician limestone of Montreal, Quebec
occurs as sills (Vard, 1993)

Objectives

- Determine specific geochemical reactions that proceed within the reservoir
- Examine propensity for self-trapping (Dawsonite precipitation)

Tools.....

- TRANS (Lichtner, 1999)
- Geochemist's Workbench (Bethke, 2002)
- Examine timing of Dawsonite precipitation
 - What is controlling this time scale?
 - How are other reactions enhancing or prohibiting Dawsonite precipitation?

Objectives


Geochemical Rxns. at high $\text{CO}_2(\text{aq})$ concentrations

pH = ~4.7


Dissolution processes

$\text{CO}_2(\text{aq}) = \sim 1$ molal

Albite dissolution




K-Feldspar dissolution



Anhydrite and Dolomite dissolution


Major Mineralization processes

****Dawsonite precipitation****



Objectives

At what conditions is Dawsonite stable?



Objectives

Parameterization of kinetic rate constants

Study	K-Feldspar (mol/cm ² -s)	SSA (cm ² /g)	Albite (mol/cm ² -s)	SSA (cm ² /g)	Dawsonite (mol/cm ² -s)	SSA (cm ² /g)
1	1.62×10^{-13}	711	3.63×10^{-13}	695	1.38×10^{-11}	849
2	1.00×10^{-16}	**	1.00×10^{-16}	**	1.00×10^{-16}	**
3	1.78×10^{-14}	**	**	**	1.00×10^{-11}	**
4	1.00×10^{-17}	300*	1.00×10^{-16}	600*	1.00×10^{-16}	300*

Notes: 300* = Specific surface area in (cm²/cm³) per bulk volume

** = no data available

1: Gauss et al. (2003)

2: Xu et al. (2002)

3: Johnson et al. (2001)

4: Stauffer et al. (2003)

Timing Study

Sensitivity Analysis to determine time scale for Dawsonite precipitation

Sensitivity Analysis: kinetic rate constants varied while holding specific surface areas constant (set to Gauss et al. (2003))


Gauss et al., 2003 parameters were used as base case

- varied K-Feldspar, Albite, and Dawsonite rate constants
- examined the timing of Dawsonite precipitation

<u>Study</u>	<u>Timing</u>
Gauss et al.	<2 yr
Xu et al.	~800 yr
Johnson et al.	~1000 yr
Stauffer et al.	~3000 yr


Timing Study

Fast kinetic rate constants: Study 1



Timing Study

Slow kinetic rate constants: Study 4



Timing Study

Controls on timing of Dawsonite precipitation

- Timing of Albite dissolution (rate constants)
 - Albite dissolution controls Al^{3+} in solution which in turn controls Dawsonite precipitation
- Kinetic rate constants of aluminosilicate minerals and of Dawsonite control the precipitation timing
- Precipitation timing varies over several orders of magnitude due to kinetic rate constant disparities

Conclusions

- Dawsonite precipitation at high $\text{CO}_2(\text{aq})$ concentrations is predicted in an arkosic sandstone reservoir
- Timing of Dawsonite precipitation varies over several orders of magnitude mainly dependent upon the kinetic rate constants of feldspathic minerals and Dawsonite

West Pearl Queen Implications

- Self-trapping may occur if separate phase CO₂ remains in the vicinity of feldspar-rich mineral zones (in the case of the West Pearl Queen reservoir, Albite)
- Dawsonite may greatly influence reservoir hydrodynamics due to porosity and permeability changes
- Sites with feldspar rich cap rocks provide the best opportunity for Dawsonite self-trapping
 - Overlying mineralogy within the West Pearl Queen reservoir is dominated by evaporitic minerals

Acknowledgements

Special Thanks to:

Brian McPherson (NMT)
Peter Lichtner (LANL)
Rajesh Pawar (LANL)
John Kaszuba (LANL)
Reid Grigg (NMT)

Gill Bond (NMT)
Ning Liu (NMT)
John Stringer (EPRI)
Dave Norman (NMT)

Research Funded by EPRI: WO9000-26 and
EP/P11940/C5920

